

JOURNAL

de Théorie des Nombres

de BORDEAUX

anciennement Séminaire de Théorie des Nombres de Bordeaux

Sinan Ünver

An Arakelov theoretic proof of the equality of conductor and discriminant

Tome 16, n° 2 (2004), p. 423-427.

<http://jtnb.cedram.org/item?id=JTNB_2004__16_2_423_0>

© Université Bordeaux 1, 2004, tous droits réservés.

L'accès aux articles de la revue « Journal de Théorie des Nombres de Bordeaux » (<http://jtnb.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://jtnb.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

An Arakelov theoretic proof of the equality of conductor and discriminant

par SINAN ÜNVER

RÉSUMÉ. Nous donnons une preuve utilisant la théorie d'Arakelov de l'égalité du conducteur et du discriminant.

ABSTRACT. We give an Arakelov theoretic proof of the equality of conductor and discriminant.

1. Introduction

Let K be a number field, \mathcal{O}_K be the ring of integers of K , and S be $\text{Spec}(\mathcal{O}_K)$. Let $f : X \rightarrow S$ be an arithmetic surface. By this we mean a regular scheme, proper and flat over S , of relative dimension one. We also assume that the generic fiber of X has genus ≥ 1 , and that X/S has geometrically connected fibers.

Let ω_X be the dualizing sheaf of X/S . The Mumford isomorphism ([Mumf], Theorem 5.10)

$$\det Rf_*(\omega_X^{\otimes 2}) \otimes K \rightarrow (\det Rf_*\omega_X)^{\otimes 13} \otimes K,$$

which is unique up to sign, gives a rational section Δ of

$$(\det Rf_*\omega_X)^{\otimes 13} \otimes (\det Rf_*(\omega_X^{\otimes 2}))^{\otimes -1}.$$

The discriminant $\Delta(X)$ of X/S is defined as the divisor of this rational section ([Saito]). If \mathfrak{p} is a closed point of S , we denote the coefficient of \mathfrak{p} in $\Delta(X)$ by $\delta_{\mathfrak{p}}$.

On the other hand X/S has an Artin conductor $\text{Art}(X)$ (cf. [Bloch]), which is similarly a divisor on S . We denote the coefficient of \mathfrak{p} in $\text{Art}(X)$ by $\text{Art}_{\mathfrak{p}}$. Let S' be the strict henselization of the complete local ring of S at \mathfrak{p} , with field of fractions K' . Let s be its special point, η be its generic point, and $\bar{\eta}$ be a geometric generic point corresponding to an algebraic closure \bar{K}' of K' . Let ℓ be a prime different from the residue characteristic

at \mathfrak{p} . Then

$$\begin{aligned} \text{Art}_{\mathfrak{p}}(X) &= \sum_{i \geq 0} (-1)^i \dim_{\mathbb{Q}_\ell} H_{\text{ét}}^i(X_{\bar{\eta}}, \mathbb{Q}_\ell) - \sum_{i \geq 0} (-1)^i \dim_{\mathbb{Q}_\ell} H_{\text{ét}}^i(X_s, \mathbb{Q}_\ell) \\ &\quad + \sum_{i \geq 0} (-1)^i \text{Sw}_{\bar{K}'/K'}(H_{\text{ét}}^i(X_{\bar{\eta}}, \mathbb{Q}_\ell)), \end{aligned}$$

where $\text{Sw}_{\bar{K}'/K'}$ denotes the Swan conductor of the Galois representation of \bar{K}'/K' . Both of these divisors are supported on the primes of bad reduction of X . We give another proof of Saito’s theorem ([Saito], Theorem 1) in the number field case.

Theorem 1. *For any closed point $\mathfrak{p} \in S$, we have $\delta_{\mathfrak{p}} = -\text{Art}_{\mathfrak{p}}$.*

Fix a Kähler metric on $X(\mathbb{C})$ invariant under complex conjugation, this gives metrics on $\Omega_{X_\nu}^1$ ’s, for each $\nu \in S(\mathbb{C})$. For a hermitian coherent sheaf \mathcal{E} , we endow $\det Rf_*(\mathcal{E})$ with its Quillen metric. The proof of the theorem has the following corollaries.

Proposition 1. *We have*

$$\begin{aligned} \deg \det Rf_*\omega_X &= \frac{1}{12} [\deg f_*(\widehat{c}_1(\omega_X)^2) + \log \text{Norm}(-\text{Art}(X))] \\ &\quad [K : \mathbb{Q}](g - 1)[2\zeta'(-1) + \zeta(-1)], \end{aligned}$$

with ζ the Riemann zeta function.

Proposition 1 is an arithmetic analogue of Noether’s formula in which $\det Rf_*\omega_X$ is endowed with the Quillen metric. Faltings [Falt] and Moret-Bailly [M-B] proved a similar formula for the Faltings metrics.

Proposition 2. *We have*

$$\frac{1}{[K : \mathbb{Q}]} \sum_{\nu \in S(\mathbb{C})} \log \|\Delta_\nu\| = 12(1 - g)[2\zeta'(-1) + \zeta(-1)],$$

where Δ_ν is the section on X_ν obtained by pulling back Δ via the map from $\text{Spec}(\mathbb{C})$ to $\text{Spec}(\mathcal{O}_K)$ that corresponds to $\nu \in S(\mathbb{C})$. In particular, the norm of the Mumford isomorphism does not depend on the metric.

2. Proof

First we prove Proposition 1. By duality ([Deligne], Lemme 1.3), $\deg \det Rf_*\omega_X = \deg \det Rf_*\mathcal{O}_X$. By the arithmetic Riemann-Roch theorem of Gillet and Soulé ([G-S], Theorem 7), we get

$$\deg \det Rf_*\mathcal{O}_X = \deg f_*(\widehat{Td}(\Omega_X^1)^{(2)}) - \frac{1}{2} \sum_{\nu \in S(\mathbb{C})} \int_{X_\nu} Td(T_{X_\nu})R(T_{X_\nu}).$$

Here Td and R are the Todd and Gillet-Soulé genera respectively, and the superscript (2) denotes the degree 2 component. Applying the definitions of these characteristic classes we obtain

$$\deg \det Rf_* \mathcal{O}_X = \frac{1}{12} \deg f_*(\widehat{c}_1(\Omega_X^1)^2 + \widehat{c}_2(\Omega_X^1)) + [K : \mathbb{Q}](g-1)[2\zeta'(-1) + \zeta(-1)].$$

Let Z denote the union of singular fibers of f , and let $c_{2,X}^Z(\Omega_X^1)$ be the localized Chern class of Ω_X^1 with support in Z (cf. [Bloch], [Fulton]). Chinburg, Pappas, and Taylor ([CPT], Proposition 3.1) prove the formula

$$\deg f_*(\widehat{c}_2(\Omega_X^1)) = \log \text{Norm}(c_{2,X}^Z(\Omega_X^1)).$$

Combining this with the fundamental formula of Bloch ([Bloch], Theorem 1)

$$-\text{Art}_{\mathfrak{p}}(X) = \deg_{\mathfrak{p}} c_{2,X}^Z(\Omega_X^1),$$

we obtain the desired formula. Note that, since $\det \Omega_X^1 = \omega_X$, $\widehat{c}_1(\Omega_X^1) = \widehat{c}_1(\omega_X)$. □

Taking degrees in the Mumford isomorphism gives

$$13 \deg \det Rf_* \omega_X = \deg \det Rf_*(\omega_X^{\otimes 2}) + \log \text{Norm}(\Delta(X)) - \sum_{\nu \in S(\mathbb{C})} \log \|\Delta_{\nu}\|.$$

The arithmetic Riemann-Roch theorem ([Falt], Theorem 3) gives

$$\deg \det Rf_*(\omega_X^{\otimes 2}) = \deg \det Rf_* \omega_X + \deg f_*(\widehat{c}_1(\omega_X)^2).$$

Therefore we get

$$(1) \quad \deg \det Rf_* \omega_X = \frac{1}{12} [\deg f_*(\widehat{c}_1(\omega_X)^2) + \log \text{Norm}(\Delta(X)) - \sum_{\nu \in S(\mathbb{C})} \log \|\Delta_{\nu}\|].$$

Subtracting (1) from the expression in the statement of Proposition 1, we obtain

$$(2) \quad \log \left(\frac{\text{Norm}(\Delta(X/S))}{\text{Norm}(-\text{Art}(X/S))} \right) = \sum_{\nu \in S(\mathbb{C})} \log \|\Delta_{\nu}\| + 12[K : \mathbb{Q}](g-1)[2\zeta'(-1) + \zeta(-1)].$$

Now X_K has semistable reduction after a finite base change K'/K . For semistable X'/S' , both $-\text{Art}_{\mathfrak{p}'}(X')$ ([Bloch]), and $\delta_{\mathfrak{p}'}(X')$ ([Falt], Theorem 6) are equal to the number of singular points in the geometric fiber over \mathfrak{p}' . Therefore $-\text{Art}_{\mathfrak{p}'} = \delta_{\mathfrak{p}'}$, and hence

$$(3) \quad \text{Norm}(\Delta(X'/S')) = \text{Norm}(-\text{Art}(X'/S')).$$

Applying this to a semistable model X' of $X \otimes_K K'$, and noting that the base change multiplies the right hand side of (2) by $[K' : K]$, we see that the right hand side of (2) is equal to zero, and hence that the equality

$$(4) \quad \text{Norm}(\Delta(X/S)) = \text{Norm}(-\text{Art}(X/S))$$

holds for X .

To prove the equality $\delta_{\mathfrak{p}} = -\text{Art}_{\mathfrak{p}}$ for an arbitrary closed point $\mathfrak{p} \in S$, we will use the following lemma.

Lemma 1. *Fix distinct closed points $\beta_1, \dots, \beta_s \in S$. For each i such that $1 \leq i \leq s$, let L_i be an extension of the completion K_i of K at β_i such that $[L_i : K_i] = n$ is independent of i . Then there exists an extension L/K such that, for each $1 \leq i \leq s$, there is only one prime γ_i of L lying over β_i , and the completion of L at γ_i is isomorphic (over K_i) to L_i .*

Proof. The proof is an application of Krasner's lemma, and the approximation lemma. Details are omitted. \square

Take $\mathfrak{p} = \beta_1$, a prime of bad reduction. Denote the remaining primes of bad reduction by β_i , $2 \leq i \leq s$. Choose extensions L_i of the local fields K_i , for all $1 \leq i \leq s$, such that L_1 is unramified over K_1 , X has semistable reduction over L_i , for $2 \leq i \leq s$, and $[L_i : K_i] = n$, for some n . Applying the lemma to this data we obtain an extension L of K . Let $T = \text{Spec}(\mathcal{O}_L)$. The curve $X \otimes_K L$ has a proper, regular model Y over T such that

- (i) $Y \otimes_T T_{\gamma_1} \simeq X \otimes_S T_{\gamma_1}$, and
- (ii) Y is semistable at γ_i , for $2 \leq i \leq s$.

Applying (4) to Y gives the equality

$$\sum_{1 \leq i \leq s} \delta_{\gamma_i} \log \text{Norm}(\gamma_i) = \sum_{1 \leq i \leq s} -\text{Art}_{\gamma_i} \log \text{Norm}(\gamma_i).$$

On the other hand because of semistability, we have $\delta_{\gamma_i} = -\text{Art}_{\gamma_i}$, for $2 \leq i \leq s$. Hence we get $\delta_{\gamma_1} = -\text{Art}_{\gamma_1}$. Since T/S is étale at γ_1 , (i) implies

$$\delta_{\mathfrak{p}} = \delta_{\gamma_1} = -\text{Art}_{\gamma_1} = -\text{Art}_{\mathfrak{p}}.$$

Acknowledgements. I would like to thank A. Abbes for his many mathematical suggestions, and for his constant encouragement. I would like to thank my adviser P. Vojta for supporting me this academic year.

References

- [Bloch] S. BLOCH, *Cycles on arithmetic schemes and Euler characteristics of curves*. Proc. of Sympos. Pure Math. **46** (1987) AMS, 421–450.
- [CPT] T. CHINBURG, G. PAPPAS, M.J. TAYLOR, *ϵ -constants and Arakelov Euler characteristics*. Preprint, (1999).
- [Deligne] P. DELIGNE, *Le déterminant de la cohomologie*. Contemp. Math. **67** (1987), 93–177.
- [Falt] G. FALTINGS, *Calculus on arithmetic surfaces*. Ann. Math. **119** (1984), 387–424.
- [Fulton] W. FULTON, *Intersection theory*. Springer-Verlag, Berlin, 1984.

- [G-S] H. GILLET, C. SOULÉ, *An arithmetic Riemann-Roch theorem*. Invent. Math. **110** (1992), 473–543.
- [M-B] L. MORET-BAILLY, *La formule de Noether pour les surfaces arithmétiques*. Invent. Math. **98** (1989), 499–509.
- [Mumf] D. MUMFORD, *Stability of projective varieties*. Enseign. Math. **23** (1977), 39–100.
- [Saito] T. SAITO, *Conductor, discriminant, and the Noether formula of arithmetic surfaces*. Duke Math. J. **57** (1988), 151–173.

Sinan ÜNVER

Department of Mathematics

University of Chicago

5734 S. University Ave.

Chicago IL 60637, USA

E-mail : sinan@math.berkeley.edu

unver@math.uchicago.edu